

Field Correlators in the Georgi-Glashow Model and Several Aspects of its String Representation

D.V.ANTONOV ^{*†}

*Institute of Theoretical and Experimental Physics,
B.Chermushkinskaya 25, 117 218, Moscow, Russia
and*

*Institut für Physik, Humboldt-Universität zu Berlin,
Invalidenstrasse 110, D-10115, Berlin, Germany*

Abstract

String representations of the Wilson loop and of the non-Abelian analog of the 't Hooft loop, defined on the string world-sheet, are studied in the Londons' limit of the 4D Georgi-Glashow model. In the first case, massive gauge bosons yield only perimeter type terms, whereas in the second case, they also interact with the string, which enables one to derive the coefficient functions parametrizing the bilocal correlator of the dual field strength tensors. The asymptotic behaviours of these functions at small and large distances are argued to be in a good agreement with the observable lattice data.

In the recent papers^{1,2}, the string representation for the 't Hooft loop average defined on the string world-sheet in the Londons' limit of the Abelian Higgs Model (AHM) has been found. In particular in Ref. 1, it has been demonstrated that the bilocal approximation to the Method of Vacuum Correlators (MVC)³ is exact in the Londons' limit, i.e. all the terms of the cumulant expansion higher than the quadratic one vanish. The value of the bilocal correlator has been derived from the propagator of the massive Kalb-Ramond field, and is in a good agreement with the present lattice data concerning this correlator in gluodynamics⁴.

In this Letter, we shall proceed to the non-Abelian case of the 4D Georgi-Glashow model and study string representations for the Wilson loop average and the non-Abelian analog of the 't Hooft loop average defined on the string world-sheet. To this end, we shall make use of the same duality transformation, which has been originally proposed in Ref. 5 and then used in Refs. 1 and 2, and earlier in 6 and 7. In this way, we shall finally also find the bilocal correlator of the dual non-Abelian field strength tensors. All the calculations will be performed in the Londons' limit of the Georgi-Glashow model. Notice however, that contrary to Ref. 7, where only the low-energy

^{*}E-mail addresses: antonov@pha2.physik.hu-berlin.de and antonov@vxitep.itep.ru

[†]Supported by Graduiertenkolleg *Elementarteilchenphysik*, Russian Fundamental Research Foundation, Grant No.96-02-19184, DFG-RFFI, Grant 436 RUS 113/309/0, and by the INTAS, Grant No.94-2851.

limit of the Georgi-Glashow model (which is in fact an Abelian theory) has been considered, in what follows we shall study this model as it is, i.e. the non-Abelian case.

We shall start with the string representation of the Wilson loop average

$$\langle W(C) \rangle = \left\langle \text{tr P exp} \left(i q \oint_C dx_\mu A_\mu^a T^a \right) \right\rangle,$$

formed by a probe external electric charge q moving along a closed contour $C : x_\mu(\tau)$, $0 \leq \tau \leq 1$, and belonging to the representation of the group $SU(2)$ with the colour “spin” J , i.e. $(T^a)^{bc}(T^a)^{cd} = J(J+1)\delta^{bd}$ (for example, $J = 1$ for the adjoint representation). To this end, we shall make use of the formula derived in Ref. 8, which enables one to replace the path ordering in the Wilson loop in the $SU(2)$ gauge theory by the integration over an auxiliary unit three-vector. Note, that this formula has already been applied for the purposes of the string representation of gluodynamics in the confining background in Ref. 9. So, the Wilson loop average we are starting with reads as follows

$$\langle W(C) \rangle = \frac{1}{\mathcal{Z}} \int \mathcal{D}A_\mu^a \mathcal{D}\Phi^a \left\langle \exp \left[\int dx \left(-\frac{1}{4} F_{\mu\nu}^2 - \frac{1}{2} |D_\mu^{ab} \Phi^b|^2 - \lambda (|\Phi^a|^2 - \eta^2)^2 + i J q j_\mu^a A_\mu^a \right) \right] \right\rangle_{\vec{n}}, \quad (1)$$

where

$$\langle \dots \rangle_{\vec{n}} \equiv \int \mathcal{D}\vec{n} (\dots) e^{i J \int_0^1 d\tau \dot{\phi} \cos \chi}$$

with the vector $\vec{n}$ being parametrized as

$$\vec{n} = (\sin \chi \cos \phi, \sin \chi \sin \phi, \cos \chi), \text{ and } j_\mu^a(x) = \int_0^1 d\tau \dot{x}_\mu(\tau) n^a(x(\tau)) \delta(x - x(\tau))$$

being its current along the contour C . The covariant derivative and the non-Abelian field strength tensor are defined as $D_\mu^{ab} = \delta^{ab} \partial_\mu - g \varepsilon^{abc} A_\mu^c$ and $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g \varepsilon^{abc} A_\mu^b A_\nu^c$ respectively.

In the Londons' limit, when the coupling constant $\lambda = +\infty$, $\Phi^a(x) = N^a \eta e^{i\theta(x)}$, where $\theta = \theta^{\text{sing.}} + \theta^{\text{reg.}}$, and $\vec{N}$ stands for a *constant* unit three-vector, whose arbitrary direction must be averaged over. In this limit, Eq. (1) reads

$$\langle W(C) \rangle = \frac{1}{\mathcal{Z}} \int \mathcal{D}A_\mu^a \mathcal{D}\theta \left\langle \exp \left[\int dx \left(-\frac{1}{4} F_{\mu\nu}^2 - \frac{\eta^2}{2} (\partial_\mu \theta)^2 - \frac{g^2 \eta^2}{3} A_\mu^2 + i J q j_\mu^a A_\mu^a \right) \right] \right\rangle_{\vec{n}}, \quad (2)$$

where the coefficient at the mass term of the field A_μ^a is just due to the averaging over the directions of the vector $\vec{N}$. Performing the duality transformation proposed in Ref. 5, we get from Eq. (2)

$$\begin{aligned} \langle W(C) \rangle &= \frac{1}{\mathcal{Z}} \int \mathcal{D}x_\mu(\xi) \mathcal{D}h_{\mu\nu} \exp \left[\int dx \left(-\frac{1}{12\eta^2} H_{\mu\nu\lambda}^2 + i\pi h_{\mu\nu} T_{\mu\nu} \right) \right] \\ &\cdot \left\langle \int \mathcal{D}A_\mu^a \exp \left[\int dx \left(-\frac{1}{4} F_{\mu\nu}^2 - \frac{g^2 \eta^2}{3} A_\mu^2 + i J q j_\mu^a A_\mu^a \right) \right] \right\rangle_{\vec{n}}, \end{aligned} \quad (3)$$

where $H_{\mu\nu\lambda} = \partial_\mu h_{\nu\lambda} + \partial_\lambda h_{\mu\nu} + \partial_\nu h_{\lambda\mu}$ stands for the strength tensor of the massless Abelian antisymmetric tensor field $h_{\mu\nu}$, $T_{\mu\nu}(x) = \int_S d\sigma_{\mu\nu}(x(\xi))\delta(x - x(\xi))$ is the vorticity tensor current, S is a closed world-sheet of the string parametrized by $x_\mu(\xi)$, and from now on, for simplicity, we shall not take into account the Jacobian emerging when one passes from the integration over $\theta^{\text{sing.}}$ to the integration over $x_\mu(\xi)$. Similarly to Ref. 5, the first line in Eq. (3) represents the interaction of a massless photon with the string world-sheet. Due to the invariance of this expression under the hypergauge transformations, $h_{\mu\nu} \rightarrow h_{\mu\nu} + \partial_\mu \Lambda_\nu - \partial_\nu \Lambda_\mu$, some gauge fixing term is needed for carrying out the integral over the field $h_{\mu\nu}$. For example, if one adds to the Lagrangian standing in the exponent in the first line of Eq. (3) the mass term $-\frac{1}{4e^2}h_{\mu\nu}^2$, the integral over the field $h_{\mu\nu}$ yields (up to a factor π^2) Eq. (2) of Ref. 10. In particular, when the regularizing photon mass $\frac{2}{e}$ is infinitesimally small, one can show that the result of the integration over the field $h_{\mu\nu}$ reads as

$$\exp\left[-\frac{e^2}{2} \oint_{\partial S} dx_\mu \oint_{\partial S} dy_\mu \frac{1}{(x-y)^2}\right],$$

(i.e. is just the one of the photodynamics case) and thus vanishes, since $\partial S = 0$. Therefore, the nonvanishing (i.e. not reducible to the boundary terms) contribution to the string effective action could arise only due to the finite photon mass, which plays the role of a hypergauge fixing parameter.

However actually, the first line in Eq. (3) drops out after the division by $\mathcal{Z}$, so that the problem of hypergauge fixing does not arise at all, and we arrive at the following expression for the Wilson loop average

$$\langle W(C) \rangle = \frac{\left\langle \int \mathcal{D}A_\mu^a \exp\left[\int dx \left(-\frac{1}{4}F_{\mu\nu}^2 - \frac{g^2\eta^2}{3}A_\mu^2 + iJqj_\mu^a A_\mu^a\right)\right] \right\rangle}{\int \mathcal{D}A_\mu^a \exp\left[-\int dx \left(\frac{1}{4}F_{\mu\nu}^2 + \frac{g^2\eta^2}{3}A_\mu^2\right)\right]}.$$

The Gaussian part of this expression reads

$$\begin{aligned} & \frac{\int \mathcal{D}A_\mu^a \exp\left[\int dx \left(-\frac{1}{4}(\partial_\mu A_\nu^a - \partial_\nu A_\mu^a)^2 - \frac{g^2\eta^2}{3}A_\mu^2 + iJqj_\mu^a A_\mu^a\right)\right]}{\int \mathcal{D}A_\mu^a \exp\left[-\int dx \left(\frac{1}{4}(\partial_\mu A_\nu^a - \partial_\nu A_\mu^a)^2 + \frac{g^2\eta^2}{3}A_\mu^2\right)\right]} = \\ & = \exp\left\{-\frac{J^2q^2}{8\pi^2} \oint_C dx_\mu n^a(x) \oint_C dy_\nu n^a(y) \frac{1}{|x-y|} \left[\delta_{\mu\nu} \left(\left(m + \frac{1}{m(x-y)^2}\right) K_1 + \frac{1}{2|x-y|} (K_0 + K_2)\right) - \right. \right. \\ & \quad \left. \left. - \frac{(x-y)_\mu(x-y)_\nu}{(x-y)^2} \left(3\left(\frac{m}{4} + \frac{1}{m(x-y)^2}\right) K_1 + \frac{3}{2|x-y|} (K_0 + K_2) + \frac{m}{4} K_3\right)\right]\right\}, \quad (4) \end{aligned}$$

where $m \equiv \sqrt{\frac{2}{3}}g\eta$ is the gauge bosons' mass, and K_i 's, $i = 0, 1, 2, 3$, stand for the Macdonald functions, whose arguments are the same, $m|x-y|$. It is worth mentioning, that during this integration one should take into account that contrary to the Abelian case, where

$$\partial_\mu j_\mu = 0 \text{ with } j_\mu(x) \equiv \int_0^1 d\tau \dot{x}_\mu(\tau) \delta(x - x(\tau)),$$

the current j_μ^a is not conserved, i.e. $\partial_\mu j_\mu^a \neq 0$.

Eq. (4) describes the interaction of the massive gauge bosons moving along the loop with each other, in the Gaussian approximation. The asymptotic behaviours of the R.H.S. of Eq. (4) at $m\sqrt{A} \ll 1$ and $m\sqrt{A} \gg 1$, where A stands for the minimal area inside the contour C , whose square root is a typical distance between two points on C , read

$$\exp \left\{ -\frac{J^2 q^2}{4\pi^2 m^2} \oint_C dx_\mu \oint_C dy_\nu \frac{1}{|x-y|^4} \left[\delta_{\mu\nu} - \frac{4(x-y)_\mu(x-y)_\nu}{(x-y)^2} \right] \right\}$$

and

$$\exp \left\{ -\frac{J^2 q^2 \sqrt{m}}{8\sqrt{2}\pi^{\frac{3}{2}}} \oint_C dx_\mu n^a(x) \oint_C dy_\nu n^a(y) \frac{e^{-m|x-y|}}{|x-y|^{\frac{3}{2}}} \left[\delta_{\mu\nu} - \frac{(x-y)_\mu(x-y)_\nu}{(x-y)^2} \right] \right\}$$

respectively.

The one-loop expression for the the Wilson loop average could be obtained upon accounting, in addition to the R.H.S. of Eq. (4), for the term resulting from the substitution of the saddle-point value of the Gaussian integral (4),

$$A_\mu^{a \text{ extr.}}(p) = \frac{iJq}{p^2 + m^2} \left[j_\mu^a(p) + \frac{p_\mu p_\nu}{m^2} j_\nu^a(p) \right]$$

into the exponent

$$\exp \left[-g\varepsilon^{abc} \int dx (\partial_\mu A_\nu^a) A_\mu^b A_\nu^c \right].$$

However, this expression again obviously does not contain any nontrivial dependence on the string world-sheet. Therefore, at least up to a one-loop approximation, in the Londons' limit, we are left only with the perimeter type contributions to the Wilson loop average.

Let us now consider the Georgi-Glashow model without external charged particles. Then the nontrivial dependence on the world-sheet S emerges in the weight factor $\langle \mathcal{F}(S) \rangle$ of the string representation for the non-Abelian analog of the 't Hooft loop average, $\int \mathcal{D}x_\mu(\xi) \langle \mathcal{F}(S) \rangle$. This weight factor has the form

$$\begin{aligned} \langle \mathcal{F}(S) \rangle = \frac{1}{\mathcal{Z}} \int |\Phi^a| \mathcal{D}|\Phi^a| \mathcal{D}A_\mu^a \mathcal{D}\theta^{\text{reg.}} \exp \left\{ - \int dx \left[\frac{1}{4} F_{\mu\nu}^2 + \frac{1}{2} |D_\mu^{ab} \Phi^b|^2 + \lambda (|\Phi^a|^2 - \eta^2)^2 + \right. \right. \\ \left. \left. + \frac{\pi}{g} \varepsilon_{\mu\nu\alpha\beta} T_{\mu\nu} F_{\alpha\beta}^a N'^a \right] \right\}, \end{aligned} \quad (5)$$

where $\vec{N}'$ is again a *constant* unit three-vector, independent of $\vec{N}$, which will drop out from the final expression for $\langle \mathcal{F}(S) \rangle$. In what follows, our aim will be to find the correlators of the dual non-Abelian field strength tensors, which could be immediately done as soon as $\langle \mathcal{F}(S) \rangle$ is known, by applying to Eq. (5) the cumulant expansion³.

In the Londons' limit, after performing the duality transformation, Eq. (5) reads

$$\langle \mathcal{F}(S) \rangle = \frac{\int \mathcal{D}A_\mu^a \exp \left\{ - \int dx \left[\frac{1}{4} F_{\mu\nu}^2 + \frac{g^2 \eta^2}{3} A_\mu^2 + \frac{\pi}{g} \varepsilon_{\mu\nu\alpha\beta} T_{\mu\nu} F_{\alpha\beta}^a N'^a \right] \right\}}{\int \mathcal{D}A_\mu^a \exp \left\{ - \int dx \left[\frac{1}{4} F_{\mu\nu}^2 + \frac{g^2 \eta^2}{3} A_\mu^2 \right] \right\}}. \quad (6)$$

Within the one-loop approximation, Eq. (6) yields

$$\begin{aligned} \langle \mathcal{F}(S) \rangle = \exp \left\{ \frac{1}{g^2} \int_S d\sigma_{\mu\nu}(x) \int_S d\sigma_{\mu\nu}(y) \left[\frac{\Lambda^4}{4} e^{-\frac{\Lambda^2(x-y)^2}{4}} - \frac{m^3 K_1(m|x-y|)}{|x-y|} \right] - \right. \\ \left. - \frac{2m}{g^2} \oint_{\partial S} dx_\mu \oint_{\partial S} dy_\mu \frac{K_1(m|x-y|)}{|x-y|} \right\}, \end{aligned} \quad (7)$$

where the first term in the exponent on the R.H.S. of Eq. (7) is due to the regularization of the delta-function with Λ standing for the UV cutoff, and the last term again vanishes, since $\partial S = 0$. Comparing now Eq. (7) with the result of the cumulant expansion of Eq. (5) and parametrizing the bilocal correlator in the adjoint representation as follows³

$$\begin{aligned} \langle \tilde{F}_{\mu\nu}^a(x) \tilde{F}_{\lambda\rho}^b(0) \rangle = \frac{\delta^{ab}}{3} \left\{ \left(\delta_{\mu\lambda} \delta_{\nu\rho} - \delta_{\mu\rho} \delta_{\nu\lambda} \right) D(x^2) + \right. \\ \left. + \frac{1}{2} \left[\partial_\mu \left(x_\lambda \delta_{\nu\rho} - x_\rho \delta_{\nu\lambda} \right) + \partial_\nu \left(x_\rho \delta_{\mu\lambda} - x_\lambda \delta_{\mu\rho} \right) \right] D_1(x^2) \right\}, \end{aligned}$$

we arrive at the following values of the functions D and D_1

$$D(x^2) = \frac{3}{2\pi^2} \left[\frac{m^3 K_1(m|x|)}{|x|} - \frac{\Lambda^4}{4} e^{-\frac{\Lambda^2 x^2}{4}} \right], \quad (8)$$

$$D_1(x^2) = \frac{3m}{\pi^2 x^2} \left[\frac{K_1(m|x|)}{|x|} + \frac{m}{2} (K_0(m|x|) + K_2(m|x|)) \right]. \quad (9)$$

The asymptotic behaviours of these functions at $|x| \ll \frac{1}{m}$ and $|x| \gg \frac{1}{m}$ read

$$D \longrightarrow \frac{3m^2}{2\pi^2 x^2}, \quad (10)$$

$$D_1 \longrightarrow \frac{6}{\pi^2 |x|^4}, \quad (11)$$

and

$$D \longrightarrow \frac{3m^4}{2\sqrt{2}\pi^{\frac{3}{2}}} \frac{e^{-m|x|}}{(m|x|)^{\frac{3}{2}}}, \quad (12)$$

$$D_1 \longrightarrow \frac{3m^4}{\sqrt{2}\pi^{\frac{3}{2}}} \frac{e^{-m|x|}}{(m|x|)^{\frac{5}{2}}} \quad (13)$$

respectively, where during the derivation of Eq. (10) we have put Λ to be of the order of m , which is true in the confining regime¹.

Thus, we conclude that the bilocal approximation to MVC is exact in the Londons' limit of the Georgi-Glashow model in the one-loop approximation, i.e. analogously to the AHM in the Londons' limit¹, all the cumulants higher than the quadratic one vanish. Coefficient functions (8) and (9) are the main result of the present Letter. Similar to the Abelian case¹, their asymptotic behaviours (10)-(13) are in a good agreement with the observable lattice data⁴ on the corresponding behaviours in gluodynamics.

Acknowledgments

The author is grateful to Professors M.I.Polikarpov and Yu.A.Simonov for the encouragement and to M.N.Chernodub for criticism and a lot of useful discussions on the string representation of AHM. He would also like to thank the Quantum Field Theory Department of the Institute of Physics of the Humboldt University of Berlin and especially Profs. D.Ebert, D.Lüst, and M.Müller-Preussker for kind hospitality.

References

1. D.V.Antonov, *hep-th/9710144*.
2. D.V.Antonov, *hep-th/9710194*.
3. H.G.Dosch, *Phys.Lett.* **190**, 177 (1987); Yu.A.Simonov, *Nucl.Phys.* **B307**, 512 (1988); H.G.Dosch and Yu.A.Simonov, *Phys.Lett.* **B205**, 339 (1988), *Z.Phys.* **C45**, 147 (1989); Yu.A.Simonov, *Nucl.Phys.* **B324**, 67 (1989), *Phys.Lett.* **B226**, 151 (1989), *Phys.Lett.* **B228**, 413 (1989), *Yad.Fiz.* **58**, 113 (1995), preprint ITEP-PH-97-4 (*hep-ph/9704301*), *hep-ph/9712248*, *hep-ph/9712250*; for a review see Yu.A.Simonov, *Yad.Fiz.* **54**, 192 (1991), *Phys.Usp.* **39**, 313 (1996).
4. A. Di Giacomo and H.Panagopoulos, *Phys.Lett.* **B285**, 133 (1992); A. Di Giacomo, E.Meggiolaro, and H.Panagopoulos, *hep-lat/9603017* (preprints IFUP-TH 12/96 and UCY-PHY-96/5) (in press in *Nucl.Phys.* **B**).
5. K.Lee, *Phys.Rev.* **D48**, 2493 (1993).
6. M.I.Polikarpov, U.-J.Wiese, and M.A.Zubkov, *Phys.Lett.* **B309**, 133 (1993); P.Orland, *Nucl.Phys.* **B428**, 221 (1994); M.Sato and S.Yahikozawa, *Nucl.Phys.* **B436**, 100 (1995); E.T.Akhmedov, M.N.Chernodub, M.I.Polikarpov, and M.A.Zubkov, *Phys.Rev.* **D53**, 2087 (1996).
7. E.T.Akhmedov, *JETP Lett.* **64**, 82 (1996).
8. D.I.Diakonov and V.Yu.Petrov, *Phys.Lett.* **B224**, 131 (1989).
9. D.V.Antonov and D.Ebert, *Mod.Phys.Lett.* **A12**, 2047 (1997).
10. D.V.Antonov, *hep-th/9707245*.